



Citrus Cultivation in Venezuela

Gustavo Fermin^{1*} • Armando Briceño¹ • Francisca Ely¹ •
Luis Cedeño² • Yani Aranguren¹ • Carle Valecillos¹

¹ Centro Jardín Botánico, Facultad de Ciencias, La Hechicera, Universidad de Los Andes, Mérida 5101, Venezuela

² Instituto de Investigaciones Agrícolas y Pecuarias, Facultad de Ciencias Forestales, Santa Rosa, Universidad de Los Andes, Mérida 5101, Venezuela

Corresponding author: * fermin@ula.ve

ABSTRACT

This short review focuses on a few aspects related to the citrus industry that have received attention in Venezuela. From a well, culturally speaking, established citrus industry, consistent efforts have been made to study the main pests and pathogens that cause diseases and economic losses in citrus orchards across Venezuela. Despite the knowledge gathered from almost a century of phytopathological studies, however, very few advances have been attained in the improvement of citrus germplasm in Venezuela. In the same way, postharvest management of the produce has not shown a successful history of accomplishments. The marginal increase of production and yield in recent times that seems to be decelerating the tendency of the last fifteen years to accumulate losses, however, have yet to demonstrate that the industry is experiencing a renewal of its cultural practices and an increase in financial and technological inputs. Increases in the consumption of fruits have also been recorded in the last few years, but consumption has not yet reached the recommended levels set by international standards. The citrus industry in Venezuela has still much to do, despite being one of the most important in the country, to contribute to food security and to offer the producers an attractive source of personal and professional satisfaction as well as financial reward.

Keywords: citriculture, fruit crops, orange

CONTENTS

INTRODUCTION.....	152
HISTORICAL BACKGROUND AND CULTURAL ADOPTION	155
Production, varieties and native Rutaceae species.....	156
PATHOGENS AND PESTS.....	157
Viroids and viruses.....	158
Bacteria.....	158
Fungi and oomycetes.....	158
Nematodes.....	159
Insects.....	159
Others: weeds, algae and mollusks.....	159
IMPROVEMENT AND PERSPECTIVES	160
CONCLUDING REMARKS	160
REFERENCES.....	161

INTRODUCTION

Aside from tasting good and being one of the most refreshing fruits, citrus are high in vitamin C content. Members of this important group of fruit plants are ubiquitously found in Venezuela. Citrus trees constitute the first perennial, woody fruit crop in Venezuela; it has also been favored as a domestic ornamental plant and may be found almost everywhere in the country, with exception of the cold, high altitude regions. Citrus fruits are among the favorite fruits for fresh consumption and juice processing in the country. In fact, they run only behind bananas and pineapple as fresh produce (Mora and Rojas 2007), and they are also the ones preferred by the juice industry, which is particularly important in a country where fresh fruits, but more importantly fruit juices, are a traditional and obligate part of the daily diet. Yet, fruit consumption in Venezuela still lags behind the recommended daily intake (Machado-Allison 2007). For the year 2004, for instance, the availability of fresh fruits in the diet of Venezuelans was 123.8 g per person, of which 16.7 corresponds to oranges (Ministerio del Poder Popular

para la Salud 2007). From all the statistics analyzed so far, it is clear that all of citrus production is consumed locally, except for a marginal exportation of 'Tahiti' lime (Laborem *et al.* 1996). Nonetheless, the perspectives of an increasing demand for food, and particularly fresh fruits, and governmental plans to increase production, might revert the negative figures associated with the citrus agroindustry in Venezuela over the last 15 years. A marginal increase in production, if sustained or improved, might permit a recovery of the production to levels similar to those reached in the early 1990s.

The citrus industry, however, faces many challenges, with pests and pathogens ranking as the main obstacle that medium to high income growers must confront. From viroids and viruses to higher eukaryotes, citrus plants in Venezuela must coexist with a myriad of potential, and actual, enemies, as reflected in **Table 1**. In this short review emphasis will be given to phytopathological aspects of citrus cultivation in Venezuela, since improvement of the germplasm currently being exploited in the country is still in its infancy – despite the fact that citrus plants have been

Table 1 Main pests, weeds and pathogens of citrus plants reported in Venezuela.*

Viroids	<i>Citrus exocortis viroid</i> <i>Citrus xyloporosis viroid</i>
Virus	<i>Citrus leprosis virus</i> <i>Citrus psorosis virus</i> <i>Citrus tristeza virus</i>
Bacteria	<i>Spiroplasma citri</i> <i>Xylella fastidiosa</i>
Fungi and oomycetes	<i>Aschersonia placenta</i> <i>Alternaria citri</i> <i>Alternaria</i> sp. <i>Ascochyta citri</i> <i>Ascochyta</i> sp. <i>Aspergillus niger</i> <i>Botryodiplodia theobromae</i> <i>Botryosphaeria rhodina</i> <i>Camarosporium</i> sp. <i>Capnodium citri</i> <i>Capnodium</i> sp. <i>Cercospora penzigii</i> (= <i>aurantia</i>) <i>Cercospora citrigrisea</i> <i>Cercospora</i> sp. <i>Chaetothyrium</i> sp. <i>Diaporthe citri</i> <i>Diplodia natalensis</i> <i>Elsinoe fawcettii</i> <i>Fusarium lateritium</i> <i>Fusarium</i> sp. <i>Gloeosporium limeticolum</i> <i>Glomerella cingulata</i> <i>Metarhizium anisopliae</i> <i>Microxyphium</i> sp. <i>Mycosphaerella citri</i> <i>Mycosphaerella horii</i> <i>Pellicularia koleroga</i> <i>Penicillium digitatum</i> <i>Penicillium italicum</i> <i>Phomopsis citri</i> <i>Phyllosticta</i> sp. <i>Physcia</i> sp. <i>Phytophthora citrophthora</i> (= <i>parasitica</i>) <i>Phytophthora palmivora</i> <i>Phytophthora nicotianae</i> <i>Phytophthora terrestris</i> <i>Rhizopus nigricans</i> <i>Septobasidium alni</i> <i>Septobasidium pseudopedicellatum</i> <i>Septoria aurantiicola</i> <i>Septoria citri</i> <i>Sphaeropsis tumefaciens</i> <i>Sphaerostilbe coccophila</i> <i>Sphaerostilbe flammea</i> <i>Sporotrichum citri</i> <i>Stenella citri-grisea</i> <i>Sirigula elegans</i> <i>Sirigula</i> sp. <i>Trametes scabrosa</i> <i>Tripospermum</i> sp. <i>Tubercularia</i> sp.
Algae	<i>Cephaleuros virescens</i> <i>Trentepohlia</i> sp. (incertae sedis)
Nematodes	<i>Criconema demani</i> <i>Criconemoides</i> sp. <i>Gracilacus aculeata</i> <i>Helicotylenchus crenacauda</i> <i>Helicotylenchus microcephalus</i> <i>Helicotylenchus multicinctus</i> <i>Hemicriconemoides coccophillus</i> <i>Hemicriconemoides communis</i> <i>Hoplolaimus seinhorsti</i> <i>Meloidogyne exigua</i> <i>Mesocriconema sphaerocephalum</i> <i>Paratrichodorus minor</i>

Table 1 (Cont.)

Nematodes	<i>Paratylenchus elachystus</i> <i>Paratylenchus minutus</i> <i>Peltamigratus holdemani</i> <i>Peltamigratus machethi</i> <i>Pratylenchus brachyurus</i> <i>Pratylenchus hexincisus</i> <i>Pratylenchus scribneri</i> <i>Pratylenchus zeae</i> <i>Radopholus similis</i> <i>Rotylenchus caudaphasmidius</i> <i>Rotylenchulus reniformis</i> <i>Scutellonema brachyurum</i> <i>Trichodorus</i> sp. <i>Tylenchorhynchus acutus</i> <i>Tylenchorhynchus annulatus</i> <i>Tylenchorhynchus capitatus</i> <i>Tylenchulus semipenetrans</i> <i>Tylenchus costatus</i> <i>Xiphinema americanum</i> <i>Xiphinema brasiliense</i> <i>Xiphinema brevicollum</i> <i>Xiphinema krugi</i> <i>Xiphinema peruvianum</i> <i>Xiphinema simile</i> <i>Xiphinema vulgare</i>
Mollusks	<i>Orthalicus maracaibensis</i> <i>Oxystiyla pulchella</i> <i>Polymita</i> sp.
Insects and arachnids	<i>Acromyrmex octospinosus</i> <i>Aleurocanthus woglumi</i> <i>Aleurothrixus floccosus</i> <i>Alabama argillacea</i> <i>Anastrepha fraterculus</i> <i>Aphis citricola</i> <i>Aphis craccivora</i> <i>Aphis gossypii</i> <i>Aphis nerii</i> <i>Aphis spiraeicola</i> <i>Atta sexdens</i> <i>Brevipalpus obovatus</i> <i>Brevipalpus phoenicis</i> <i>Chrysomphalus aonidum</i> <i>Coccus viridis</i> <i>Cratosomus punctulatus</i> <i>Crematogaster</i> spp. <i>Dialeurodes citri</i> <i>Diaphorina citri</i> <i>Diaprepes abbreviatus</i> <i>Dysdercus maurus</i> <i>Eutetranychus banksi</i> <i>Epitragus fuscus</i> <i>Frankliniella</i> sp. <i>Glyptoscelis fascicularis</i> <i>Gonodonta clotilda</i> <i>Gonodonta nutria</i> <i>Gonodonta pyrigo</i> <i>Gymnandrosoma auriantanum</i> <i>Heliothrips haemorrhoidalis</i> <i>Icerya purchasi</i> <i>Lampetis cacica</i> <i>Lepidosaphes beckii</i> <i>Lepidosaphes gloverti</i> <i>Litostylus juvenicus</i> <i>Oligonychus peruvianus</i> <i>Orthezia</i> sp. <i>Panonychus citri</i> <i>Papilio anchisiades</i> <i>Papilio thoas</i> <i>Parabemisia myricae</i> <i>Paraleyrodes</i> sp. <i>Parlatoria ziziphi</i> <i>Phobetrus hipparchia</i> <i>Phyllocoptruta oleivora</i>

Table 1 (Cont.)

Insects and arachnids	<i>Phyllocnistis citrella</i> <i>Pinnaspis aspidistrae</i> <i>Platynota</i> sp. <i>Platytlelus costalis</i> <i>Polyphagotarsenonius latus</i> <i>Pseudococcus</i> sp. <i>Saissetia hemisphaerica</i> <i>Saissetia nigra</i> <i>Selenaspis articulatus</i> <i>Solenopsis geminata</i> <i>Tetranychus trinidadensis</i> <i>Tetranychus tumidus</i> <i>Toxoptera aurantii</i> <i>Toxoptera citricida</i> <i>Trigona amalthea</i> <i>Trigona trinidadensis</i> <i>Trigona</i> spp. <i>Unaspis citri</i>
Weeds	
Dicots	<i>Acmeilla radicans</i> <i>Clibadium surinamense</i> <i>Cuphea melvilla</i> <i>Dioeclea guianensis</i> <i>Fleischmannia microstemon</i> <i>Galinsoga quadriradiata</i> <i>Gamochaeta americana</i> <i>Hyptis atrorubens</i> <i>Irlbachia alata</i> <i>Jacobina boliviensis</i> <i>Justicia pectoralis</i> <i>Mikania micrantha</i> <i>Odonellia hirtiflora</i> <i>Physalis angulata</i> <i>Prestonia acutifolia</i> <i>Ruellia geminiflora</i> <i>Salvia tiliifolia</i> <i>Sipanea pratensis</i> <i>Solanum americanum</i> <i>Vismia baccifera</i>
Monocots	<i>Cipura paludosa</i> <i>Trimezia martinicensis</i>
Other plants (parasitic)	<i>Langsdorfia</i> sp. <i>Phoradendron</i> sp.

*The bibliographic source for the organisms listed here is multiple, and they are available upon request. However, we recommend the following references that have shown to provide the most useful and complete information: for general references, Fundación Shell (1966), Ordozgoity *et al.* (1971), Díaz and Salas (1973), and Junta del Acuerdo de Cartagena (1996); for phytopathogenic nematodes, Crozzoli (2002); fungi and Chromista, Chardon and Toro (1934), Müller (1940), Spaulding (1961), Dennis (1970), Uriaga (1986), Rondón (1990, 1992), Erwin and Ribeiro (1996), Pretorius *et al.* (2003) and Farr *et al.* (2008); weeds, especially from Monagas state Lárez (2007a, 2007b); and insects, Fernández *et al.* (1957), Arnal *et al.* (1993), Briceño (2007) and Cermeli (2007).

amongst us for at least 400 years –, and plant protection is the aspect of citrus production that has received the utmost attention in the country.

Besides their natural enemies, as mentioned before, citrus cultivation in Venezuela has been historically characterized by some positive, but also negative factors that conspire against a successful degree of accomplishment. Fruit cultivation has a long tradition in the country as a cultural practice that has received careful attention from phytopathologists, entomologists and agronomists. Citrus cultivation in Venezuela, however, is characterized by: 1) being almost exclusively devoted to cope with local demands, although insufficiently; 2) a lack of rational plans to improve or develop cultivars adapted to our conditions; 3) low yield values mostly due to inappropriate cultural practices, mismanagement of pests and pathogens, and an almost non-existent improvement of the germplasm employed to sustain both fresh consumption and industrial processing; 4) currently used post-harvest practices (inadequate selection,

improper transportation) that render a poor performance; 5) being severely affected by a low level of technical support and an almost inexistent evaluation of production means and practices; 6) absence of a concerted relation between research and production; 7) a consuetudinary instability of official support policies, particularly demonstrated by erratic measures regarding agriculture broadly speaking, and citriculture in particular; 8) in general, quality of the products is low and phenotypes are heterogeneous.

Production of combined citrus is mainly managed in the central-northern states of the country (Yaracuy, Carabobo and Aragua), Monagas and Sucre to the east, and Táchira and Zulia in western Venezuela (Fig. 1), but minor producing units are also present in the rest of the country, with exception of south to Orinoco River. Although other states of the country apparently also show appropriate conditions for citrus production (Rudolph 1944), diverse economical, social and political reasons have precluded the extensive use of citrus species in areas recommended for their cultivation. For a comprehensive agroecological zone description of orange production in Venezuela, however, we encourage the reader to consult an excellent monograph by Benaccio *et al.* (1985) in which the authors describe the physical and climatic characteristics of promising areas where citrus might be cultivated, state by state.

It has already been mentioned that citrus species have been present in Venezuela for almost 400 years; but commercial production at great scale, however, is a recent development that can be traced back to the 1960s. In this regard citrus cultivation in Venezuela is still a nascent industry that, when considering its short life, has been somewhat successful. Attempts to increase yield, for example, has responded to the need of increasing value and financial return, but also, contradictorily, to pest management. When citrus plants succumbed to tristeza in the early 1980's, not only more appropriate rootstocks were used to substitute susceptible ones, but crop density also increased, and hence, production – as demonstrated by the golden years of citrus production during the next decade. Effective density from 156-204 plants/ha increased to 319-408, or even to 1283 plants/ha when using dwarfing germplasm (Citrumelo 'Swingle'), as reported by Avilán and Ruiz (1999). Increase in crop density, as well as the application of effective phytosanitary measures, should help producers sustain and increase production; the latter, however, has always been the weakest point of Venezuelan fruitculture, although successful cases of biological control, however, will be presented later. On the other hand, it has been shown that plants bear fruits during 7-11 months, and studies to determine best time of harvest, under Venezuela conditions, have also been performed (Laborem *et al.* 1993). Despite these efforts to study some factors related to crop management (crop density, plant fertilization, time of harvest, etc.), and to apply them to the limited cases in which producers are willing to take the risk, citriculture in Venezuela is still characterized by an almost permanent low productivity.

Low yield (less than half of what is potentially achievable) is attributed to the low technical input of the agroindustry, as well as to the reasons mentioned in preceding paragraphs. If the citrus industry, then, is to contribute to the long aspired goal of food security in the country, at least four objectives must be pursued: 1) Careful selection of a germplasm that will allow for extensive studies related to improvement, exploitation, production, processing and diffusion among consumers in order to offer to vulnerable groups an easy access to macro, micronutrients and other compounds (e.g., vitamins and antioxidants) that might positively impact their health and wellbeing; 2) Complete assessment of the totality of chemicals in citrus plants that might be useful to attain the previous goal of germplasm improvement. These include, of course, historical uses of citrus chemicals that, perhaps due to lack of study, have not been exploited adequately, but that might contribute to an increase in the possibilities of a wider use and consumption of different citrus species. In the long term, the identifica-




Fig. 1 Data of orange production in Venezuela. (A) Map of harvested area for oranges, shown by state. (B) Map of oranges production, shown by state. (C) Area devoted to the cultivation of oranges (in Ha), as well as yield (in Kg/Ha) during the last 15 years. (D) Orange production in Venezuela during the last 15 years. Maps were generated by the FAO on-line interactive service (<http://www.fao.org/landandwater/agll/agromaps/interactive/page.jsp>), and data for graphs C and D were taken from the statistical database of Fedeaagro (<http://www.fedeaagro.org/>).

tion of major germplasm, as well as the weak points that need a more conscious action, will indicate which genetic features, biosynthetic pathways, and resistance performance determinants need to be genetically altered in order to improve and strengthen the citrus industry in Venezuela; 3) A sustained increase in the use of different techniques and strategies based on molecular markers to characterize old and new varieties and assess their purity, as well as a detailed description and catalogue of the main pathogens and their population structures. Additionally, the mastering of molecular knowledge and techniques will allow engineering changes in composition and physiology of the produce during maturation, harvesting, transportation, either by classical genetic improvement practices or by transformation by different means; and 4) a concerted action between researchers, producers and governmental representatives which will permit to establish a coherent set of regulations concerning all aspects related to the citrus industry, including germplasm manipulation and phytosanitary measures.

HISTORICAL BACKGROUND AND CULTURAL ADOPTION

All first accounts from the conquerors that first visited and described the newly discovered country (1498, and later) indicate that the diet of native Venezuelans was rich in fruits that included guava (*Psidium guajava*), papaya (*Carica papaya*), diverse Annonaceae, a few of the genus *Spondias*, several palms and pineapples, as well as other minor fruits (Patiño 1963; Cumill *et al.* 2007). Although the first conquerors found difficulty in swallowing most of the native fruits,

other fruits, like pineapple, were quickly adopted. Nonetheless, the Spaniards introduced, as soon as they could, those fruits that were familiar to their diet. Among the first fruits introduced by Spaniards and Portuguese to tropical America were oranges, along with other *Citrus* species that included lemon and lime. The entrance of citrus trees to Venezuela, in particular, took place at different times. At the beginning, it was part of the first attempt to implant in the newly discovered land the agricultural goods and means they knew in Spain; years later, some citrus fruits were introduced again by slaves brought to America by the Europeans. In either case, the origin of citrus plants in Venezuela is the same: from Asia, by way of adoption in Europe and Africa, and their subsequent cultural implantation in our land (Patiño 1969).

First historical references, from different sources, to citrus fruits adopted early in Venezuela include *C. aurantiifolia* (1573), *C. medica* (1579), *C. decumana* (1578) and *C. aurantium* (1578). Other citrus plants, such as *C. grandis* and *C. reticulata*, arrived later (XIX century), but were readily accepted, especially since the first adopted citrus created a culture of sour fruit consumption. However, it is likely that citrus cultivars were introduced earlier, since by 1535 some varieties were already reported in the Caribbean (Vila 1981). In fact, according to Patiño (1969), citing different sources, Columbus brought seeds of orange, lime, cider and lemon as early as 1493, during his second journey to America.

Citrus plants are not only an integral part of the modern Venezuelan agriculture; these were readily adopted from the very moment natives were acquainted and used to their con-

sumption, even at present, by diverse ethnic groups that maintain their indigenous identity. Among the Hiwi in Venezuela, for example, *C. aurantium* is one of the common seasonal fruits included in a diet where mangoes, and other less well-known fruits, can also be found (Hurtado and Hill Kim 1990). None of these two fruits are autochthonous, and yet, they constitute an integral part of the nutritional culture of indigenous people, that blend well with their traditional background derived from nature and their ancestors. Among the Yanomami, inhabiting the Brazilian and Venezuelan Amazon, on the other hand, several autochthonous Rutaceae, although not *Citrus*, are used for different medical purposes (Milliken and Albert 1997). Finally, in various agriculture-based societies, mainly in the Andean states, citrus was also easily adopted, among other crops that enriched the cultural practices of the conquered natives (Clawson and Raymond 1982).

Citrus fruits have a wide variety of traditional uses worldwide, and very recently have gained additional public interest due to their high content in ascorbic acid, flavonoids, lycopenes, carotenes and lemonoids (Ferguson 2002). Although the fruits are mainly consumed as fresh fruits in Venezuela, juices or desserts are prepared domestically or industrially. Citrus fruits and different organs of the plant are used in ethnomedicine, as any Venezuelan is familiar with at least one of their various beneficial properties or uses, particularly in the case of oranges and lemons. Customarily, Venezuelans believe in the health-promoting effects of vitamin C, which they believe is mainly found in orange and lemon; the latter being preferred for the treatment or alleviation of cold symptoms, especially if mixed with honey (Vit *et al.* 2006). However, the studies concerning the medical uses of Rutaceae in Venezuela are scarce (Játem *et al.* 1997), which is surprising considering the amount of people familiar with the use and commerce of herbs and plant parts used in popular pharmacopoeia, yet they are unaware of the exact health promoting properties of citrus plants.

Leaves, flowers and fruits are used, alone or in combination with other natural products, in beauty treatments and in the relief of ailments of the respiratory, digestive, endocrine, cardiovascular and muscle-skeleton systems; also as analgesic, and for the cure of infectious and parasite-derived diseases (Jones *et al.* 1964; Keshava 1985; Caballero 1995; CONAPLAMED 2000; Bermúdez and Velásquez 2002; Gil and Carmona 2003; Lárez 2004; Carrillo-Rosario and Moreno 2006), or as disinfectant in the control of poultry coccidiosis (Tamasaukas *et al.* 1997).

C. aurantiifolia and *C. limon* are used as respiratory antiseptic, gastric protectant, antiemetic, diuretic, spasmolytic, antidepressant, antifungal, wound healer, to improve digestion and blood circulation, and for the treatment of diarrhea, the alleviation of cold and coughing, and for insect bites (Jones *et al.* 1964; Keshava 1985; Hoyos 1994; CONAPLAMED 2000; Bermúdez and Velásquez 2002; Carrillo-Rosario and Moreno 2006). In addition, they are also used in beauty treatments of skin and hair (Keshava 1985). *C. aurantiifolia* has also been used as an antihemorrhagic to treat snake bites (Lárez 2004).

C. sinensis is employed in the treatment of asthma, coughing, hypertension, spasms, pain, diabetes, and also as an alternative for the management of obesity, malaria and some other hepatic and cardiac problems (Hoyos 1994; Gil and Carmona 2003), and as a febrifuge and antilflatulent (Lárez 2004). For most of the 1930s, orange leaves mixed with baking soda was the only dental care product available in eastern rural Venezuela (Muñoz de Fermin, pers. comm.). *C. reticulata* and *C. reshni* are preferred for the health and care of the skin, control of hormonal disorders, and as a diuretic (Keshava 1985; Hoyos 1994). *C. medica*, on the other hand, has been reported as an efficient alternative as antiemetic, antiseptic, and vermifuge (Hoyos 1994), while *C. x paradisi* is also popular for the control of hypertension (Lans 2006).

Pulp and flour, made of seeds and peel as subproducts

of the juice industry, are used in the concentrate food for bovines, poultry and fishes (Pérez *et al.* 1975; Moreno *et al.* 2006), whilst essential oils from *C. reticulata* have showed an efficient antimicrobial activity (Martínez *et al.* 2003), and those derived from lemon as an effective insect repellent (Rojas and Scorza 1991). Some varieties of citrus and a few other Rutaceae are popular as ornamental plants in domestic gardens, or as live fences, including *C. aurantium*, *C. sinensis*, *C. madurensis*, *C. lansium*, *Fortunella margarita* and *Triphasia trifolia* (Serpa 1965; Hoyos 1994).

Finally, of the autochthonous Rutaceae, some are preferred for wood, while others, like *Pilocarpus goudotianus*, are popular for its renowned properties in the treatment of glaucoma (Hermoso and Escala 2002). Another non-indigenous Rutaceae, *Ruta graveolens*, or common rue, is employed as a carminative and for menstrual pains (Lans 2007).

Citrus was not only well accepted after their introduction; they are now an integral part of Venezuelan consumption and pharmacopoeia, as they blend well with the fruits of the neotropics, forming a rich assortment of fruits uncommon to many regions of the world.

Production, varieties and native Rutaceae species

The Rutaceae family is a group of mostly woody, very fragrant plants, comprised by 155 genera and 930 species, both tropical and subtropical (Judd *et al.* 2002). The best known and representative member of this family is the genus *Citrus*, whose species are of great economic importance due to their fruits and essence oils (Mabberley 2000). Citrus are original from tropical and subtropical Asia, specifically China (Gmitter and Hu 1990), although the taxonomy of the group is very confusing (Scora 1975). However, molecular markers are helping elucidate relationships among different *Citrus* species (for an example in oranges and lemon see Moore 2001). In Venezuela there are 25 native genera of Rutaceae plus one exotic genus (*Citrus*), with 84 native or naturalized species, including 16 that are endemic (Ricardi 1991; Hokche *et al.* 2008). Among these endemics are the genera, *Apocaulon*, *Galipea*, *Pilocarpus*, *Rutaneblina*, and *Zanthoxylum* (Taylor 1989; Reynel 1995). *Pilocarpus* and *Zanthoxylum* have some potential for their medical uses and for their wood, respectively (Patiño 1982).

Citrus is the most important plants from the Rutaceae family in Venezuela. Currently, the main producing areas comprise the states of Carabobo, Yaracuy, Aragua and Miranda, in the central region of the country, the states Monagas and Sucre in the east, and Táchira (Fig. 2) and Zulia in the west (Moreno *et al.* 2006), as shown in Fig. 1. However, it is fair to say that citrus became so well adapted and was adopted from the very beginning of the colonization that they ended up being ubiquitous in the country, to the point that the nascent colonial industry of cattle considered guava and citrus trees as real pests (Patiño 1969).

As mentioned previously, citrus fruits comprise the second group of fruits produced in the country, which employs 30–40,000 Ha for an annual production that oscillates around the half million tons per year (Fig. 1). The most important citrus varieties, in terms of production, are oranges (60.05%), tangerines (26.72%), limes (11.91%), and others including grapefruits. Yield has been estimated at 13.3, 16.4, 16.3 and 9.6 TM.Ha⁻¹, respectively. In general, there is an ample variation in crop density, with a predominance of the low ones, with modern irrigation practices limited to few orchards, a heterogeneous use of fertilizers and infrequent weeding, either manual or mechanical.

There is considerable controversy and confusion regarding the taxonomic designation of the species and varieties of citrus cultivated in Venezuela (Serpa 1965). For this reason, aside from the lack of an organized citrus industry and deficient germplasm conservation in Venezuela, the actual and current genetic citrus resources present are not well known. However, in Venezuela diverse species of *Citrus* are cultivated: *C. sinensis*, *C. aurantium* L., *C. maxima*, *C. reti-*




Fig. 2 Citrus cultivation in Táchira, western Venezuela. (A) A tangerine tree from an orchard close to La Tendida, in one of the most efficient citrus producing areas of the Táchira state. (B) Flower of *C. latifolia*, at a different orchard in the same area. (C) Tangelo fruits grown nearby La Tendida. (D) Typical symptoms of tristeza, in an adult orange tree. (E) Symptoms of (treated) gummosis in orange. (F) Symptoms of an unidentified virosis in tangelos.

culata, *C. x paradisi*, *C. grandis*, *C. limon*, *C. aurantiifolia*, *C. taiwanica*, *C. sunki*, *C. latifolia*, *C. reshni*, *C. jambhiri*, *C. volkameriana*, *C. madurensis*, *C. limetta*, and *C. medica* (Serpa 1965; Hoyos 1994; Quijada *et al.* 2002). In addition, there are non-*Citrus* citrus plants of other related genera, such as *Fortunella margarita*, *F. japonica*, *Poncirus trifoliata*, *Clausena lansium* and *Triphasia trifolia* (Serpa 1965; Hoyos 1994). Despite the lack of a historical register concerning the many occasions citrus have been re-introduced in Venezuela, during the first half of the twentieth century the favorite donor countries of citrus germplasm were Puerto Rico and Florida, especially for research purposes (Pacheco 2003).

Due to crop decline caused by phytosanitary, cultural and environmental problems, it has been customary to use rootstock grafting which has permitted, with a varying degree of success, to save the production by means of increased use of plants that are more resistant, better adapted or more productive (Rondón *et al.* 1993; Monteverde *et al.* 1996; Laborem *et al.* 1998). Nonetheless, despite the advantages that this germplasm offers, new problems have emerged that require the search and development of new species and varieties able to cope with pressing challenges (Rondón *et al.* 1993). By the year 1982 and later, preferred rootstock for commercial production in Venezuela included, for example, 'Citrange Troyer', 'Citrange Carrizo', 'Citrumelo Swingle', 'Mandarina Cleopatra', 'Limón Volkameriano', *Citrus taiwanica* and 'naranja criollo', aimed at dealing with main citrus diseases that included tristeza, exocortis, xyloporosis and gummosis. There has also been a growing interest dealing with vigor, drought resistance, productivity and fruit quality (Leal and Trujillo 1970); however, no consistent effort has been made to improve germplasm regarding these factors.

C. aurantium is known in Venezuela as sour orange, box orange or 'naranja cochinerá', and is represented by three subspecies, *amara*, *bergamia* and *myrtifolia*. The most common of these subspecies is *amara*, which is extensively used as a rootstock (Serpa 1965; Monteverde *et al.* 1996).

C. sinensis, or sweet orange, is probably the most popular of all citrus in Venezuela. Many varieties of early, medium or late maturation have been cultivated, like 'Valencia', 'California' (Washington Navel), 'Pineapple', 'Parson Brown', 'Mediterránea dulce', 'Luegim-Gong', 'Puerto Rico', 'Frost Valencia', 'Cutter Valencia', 'Campbell Valencia', 'Frost Navel', 'D. Joao', 'Rotuna Island', 'Salustiana', 'Hamlin', 'Criollo Montero', 'Natal', 'Pera' and 'Parson Brown' (Serpa 1965; Monteverde *et al.* 1996; Laborem *et al.* 1998).

Tangerines are represented in the country by *C. reticulata* and *C. reshni*. Among the cultivated varieties, 'Willow Leaf', 'Algeria', 'Dancy', 'Cleopatra', 'Nova', 'Cravo', 'Kinnow', 'Kara', 'Poncan', 'Parson Special', and the clementines 'Fina', 'Marisol' and 'Oroval', are the most popular (Laborem *et al.* 1998; Serpa 1965). On the other hand, grapefruit *C. x paradisi* is available in varieties of white or pink flesh, and the most cultivated varieties include 'Marsh', 'Foster' Reed G.F.T., 'Reiking Pomelo', 'Duncan', 'Star Ruby' and 'Thompson' (Serpa 1965; Avilán *et al.* 1982; Laborem *et al.* 1996).

Varieties of *C. limon* are few and include 'Eureka', 'Verna', 'Lisboa', 'Villafranca' and 'Génov' (Cañizares and Salcedo 1994). This species, however, has lost commercial value since it has been substituted by varieties of *C. aurantiifolia*, *C. latifolia*, *C. volkameriana* and *C. jambhiri* (Valbuena 1996; Ojeda *et al.* 1998; Quijada *et al.* 2002).

The hybrids cultivated in the country include tangors (*C. sinensis* x *C. reticulata*), with at least three different varieties, 'Ortanique', 'Temple' and 'Minneola' (Laborem *et al.* 1998), and citranges (*C. sinensis* x *P. trifoliata*), with the varieties 'Carrizo', 'Troyer' and 'Uvalde' (Reyes and Ruiz 1984). Other hybrids cultivated in Venezuela include *C. reticulata* x *C. paradisi*, *C. reticulata* x *Fortunella* and *P. trifoliata* x *C. paradisi* (Reyes and Ruiz 1984; Serpa 1965). *C. grandis*, *C. maxima*, *C. medica*, *F. margarita*, *F. japonica*, *P. trifoliata*, *C. lansium* and *T. trifolia*, are species that are not extensively cultivated, yet can be found in small orchards and gardens (Hoyos 1994; Serpa 1965).

Finally, many rootstocks have been assayed, of which the most popular ones are those that show a certain degree of resistance against pathogens, like *C. reshni* (rootstock 'Cleopatra'), *C. aurantium* or *C. sinensis* x *P. trifoliata*. Scions are very diverse, but the most commonly found are those that are commercially attractive, such as the orange varieties 'Valencia', 'California' (Washington Navel) and the lemon *C. latifolia* (Serpa 1965; Reyes and Ruiz 1984; Valbuena 1996; Laborem *et al.* 1998; Quijada *et al.* 2002).

PATHOGENS AND PESTS

A diseased plant is incapable of completing its life cycle, due to unfavorable environmental factors, a compromised physiology or due to pathogenic agents. Fruit crops do not escape this situation, and perhaps the high sugar content of their fruits makes them an easy and attractive target of different pathogens. The list of enemies of commercially cultivated citrus is long. The list shown in Table 1 clearly reveals the broad diversity of pathogens and pests of citrus

present in Venezuelan orchards. Some of them are restricted to specific areas of the country, whilst others are found ubiquitously wherever citrus species are grown. The main challenging pathogens and pests of citrus plants in Venezuela will be shortly described below; however, some readers might like to consult the references given in **Table 1** for more comprehensive information. Diseases of unknown etiology will not be covered here, e.g. citrus blight (EPPO 2004), unless the nature of the agent is at least suspected (viruses, for example).

Viroids and viruses

Probably the most prevalent disease of citrus is tristeza, caused by the *Citrus tristeza virus* (CTV, Closteroviridae), which causes tree decline, stem-pitting and yellowing of the seedlings. At least two different genotypes, derived from the analysis of three different isolates of CTV, are present in the country as molecularly established by Hilf *et al.* (2005), yet a more comprehensive analysis is needed to characterize the population structure of the virus present in the country. The disease was first described, or identified, in Venezuela by Ciferri in 1950, and was later corroborated by Knorr in 1960 in lemon trees, *C. aurantiifolia*, and lemon 'Meyer' *C. meyeri* (Ciferri 1950; Knorr *et al.* 1960, cited by Rangel *et al.* 2000). In 1983, two conditions in Venezuela became highly conducive for the virus disease: first, that 90% of orange orchards consisted of a highly susceptible rootstock-scion combination (sweet orange/sour orange); and second, the presence of the efficient virus vector *Toxoptera citricida* (Monteverde 1983), which was first identified in the country in 1976 and was widespread by 1979 (Mendt 1992). It has been argued that CTV has caused the loss of 10 million citrus trees in Venezuela (Anonymous source, 1995, cited by Gianessi *et al.* 2002), yet we have found no references or statistics that support this assumption. Finally, cross protection has been claimed to be (successfully) used in Venezuela to control tristeza (Ochoa *et al.* 1993), yet no further information to sustain this assertion has been found.

Another viral disease of importance in Venezuela is citrus leprosis, which became a phytosanitary problem in the early and mid 1990s when prevalence of the disease increased dramatically. Restrictions imposed by neighboring countries on the importation of fresh fruits produced in Venezuela is an example of the many challenges the citrus industry faces in this country, especially when phytosanitary measures are less than adequate. The disease is caused by the *Citrus leprosis virus* (EPPO 2007), a rhabdovirus, which in Venezuela seems to be transmitted by the mites, *Brevipalpus obovatus* and *B. phoenicis* (Knorr and Denmark 1970; Gamsey *et al.* 1988). Although a protocol for the phytosanitary certification of citrus, based on RT-PCR using the viral RNA, has been standardized in the country (Rangel *et al.* 2006, unpublished), no official adoption has been followed.

Amongst the diseases attributed to undescribed viruses is *impietratura* (tissue hardening). During the early stages of infection, the disease can be detected by the appearance of very small fruits. Diseased fruits are extremely hard, gum pockets are formed on the fruit albedo, and over the fruit itself hard protuberances are also evident; production is diminished and immature fruits frequently fall (Ortega 1986). The former was reported for the first time in 1970 by Rondón on sweet orange (*C. sinensis*) varieties 'Valencia' and 'Puerto Rico' (Rondón *et al.* 1970; Monteverde *et al.* 1992). Another disease whose etiological agent is an undescribed virus, reported for the first time by Monteverde *et al.* in 1983 (Monteverde *et al.* 1992) in Venezuela, is concave gum, mainly on sweet orange. This pathogen leads to the production of scales on the cortex of the trunk and main branches, accompanied by chlorosis and diminished production (Ortega 1986).

So far, only two diseases caused by viroids have been described in Venezuela. The first, *exocortis*, is caused by *Citrus exocortis viroid* (Pospiviroid), which mainly attacks

lime 'Tahiti', leading to a reduction in trees growth. Also, affected plants produce gum exudates, exhibit longitudinal cracks and a reduced production (Ortega 1986). The disease was reported by Estrada and Malaguti in 1972 (Monteverde *et al.* 1981, 1992). The second, xylosporosis, or cachexia, is caused by the *Citrus cachexia viroid* (Hostuviroid) that mainly attacks tangerines. Diseased plants show pits on the trunk, gum pockets, yellowing of leaves and reduced production (Ortega 1986). It was reported by Monteverde *et al.* in 1985.

Bacteria

Few articles and scientific printed material have been published regarding the presence of phytopathogenic bacteria on citrus cultivated in Venezuela. In some cases, the presence of a vector has prompted some researchers to warn on the potential presence of the harbored bacteria; yet no conclusive evidence has been presented of the bacteria themselves. For instance, *Candidatus liberibacter americanus*, vectored by the psyllid, *Diaphorina citri*, (Cermeli *et al.* 2000; Halbert and Nuñez 2004) and causal agent of huanglongbing, has been actively pursued although not even preliminary data has currently been published that demonstrate its presence in Venezuela. Truly identified bacteria include *Xylella fastidiosa*, the causal agent of citrus variegated chlorosis (Hernandez *et al.* 1993), and potentially *Spiroplasma citri* (EPPO 2003a). Venezuela has been acknowledged to be free from all strains of *Xanthomonas* pathogenic to citrus (Kyprianou 2006), but some controversy exists, since several Venezuelan researchers have suggested that *X. citri* may be present in some of the citrus orchards. Yet again, there is no published data to support these claims.

Fungi and oomycetes

Amongst the most important citrus diseases reported in the phytopathological literature in Venezuela are those caused by fungi and oomycetes. Damage caused by, and the presence of the fungi, *Rhizoctonia solani*, *Pythium* sp., *Fusarium* sp., and the oomycete, *Phytophthora citrophthora* (= *parasitica*), are frequently observed in seedling beds; while in nurseries, diseases like wilting caused by *P. parasitica* and *Fusarium* sp., leaf spot by *Cercospora penzigii* (= *aurantia*) (Díaz and Salas 1973; Urriaga 1986), scab by *Sphaeceloma fawcettii*, anthracnose by *Colletotrichum gloeosporioides*; areolate leaf spot by *Pellicularia filamentosa* and diplodia gummosis and stem-end rot by *Botryodiplodia theobromae* (Rondón 1992), are fairly common. The latter, reported in 1992 by Cedeño and Palacios-Prü, is also considered the causal agent of cortical lesions and gummosis in lemon (*C. aurantiifolia*) and orange (*C. sinensis*).

In adult plants, premature drop of flowers and young fruits, especially in *C. latifolia*, are induced by fungi like *C. gloeosporioides*, *C. acutatum* and *Rhizopus* sp. (Rivas *et al.* 2006). These three pathogens have been consistently isolated from infected flowers and fruits. Another disease common in adult trees, includes leaf spot, which leads to foliage drop as observed in *C. aurantium* and lemon (Rondón 1967, cited by Contreras and Rondón 1980). Controlled infection assays performed on *C. jambhiri*, *C. aurantiifolia*, *C. volkameriana*, *C. reshni* and *P. trifoliata* x *C. sinensis*, Contreras and Rondón (1980) demonstrated that the causal agents of the disease were *Alternaria citri* and *C. gloeosporioides*; the former as a primary agent of the disease, and the latter as a synergistic factor of infection.

According to Rondón (1990), amongst the frequent and economically important diseases found in citrus orchards are gummosis or root rot, caused by *P. citrophthora*, die-back by *B. theobromae*, leaf-spot by *C. penzigii* and *C. citrigrisea*, anthracnose caused by *C. gloeosporioides*, and fumagine by *Capnodium citri*. Another disease affecting leaves and fruits is the greasy spot (*Mycosphaerella citri*, as reported by Rondón 1990 and Pretorius *et al.* 2003). Although it is not known when the pathogen appeared in the

country for the first time, it has been clearly established that it is present in citrus production areas characterized by an elevated relative humidity, abundant rain and temperatures ranging from 24 to 30°C, especially in the case of sweet orange, grapefruit and tangerine orchards. Another citrus disease of certain importance in Venezuela is citrus scab, caused by the fungus *Sphaceloma fawcettii*, the anamorph of *Elsinoe fawcettii*, which can infect true lemons, and to a lesser extent, tangerines and grapefruits (Urriaga 1986). The disease is characterized by protruding, corky lesions, which in most cases can deform or diminish the size of affected plant organs. Finally, *Septoria citri* has also been reported in Venezuela, which is reputedly the causal agent of lesions in the form of concentric rings (the fungus fruiting bodies), or septoria spot, on diverse varieties of orange, lemon and grapefruit (Dennis 1970; Rondón 1992).

Nematodes

There are many nematodes that are considered to be etiological agents of important diseases in fruit trees, particularly of citrus, worldwide (Baines *et al.* 1978; Cohn 1972; Ducharme 1969, cited by Petit 1990a). Besides fungi, nematodes are the best studied plant pathogens in Venezuela, particularly those affecting fruit trees.

First reports of nematodes on citrus in Venezuela date back to 1955, when they were collected in orchards located in the central part of the country at La Victoria, Aragua state (McBeth 1956, cited by Petit 1990a). In the late 1960s, Yopez and Meredith (1970) report on the most important nematodes that affect fruit trees including citrus, Musaceae, mangoes, avocado, pineapple and papaya. Years later, based on a systematic sampling on the main citrus producing areas of the country Petit (1990a, 1990b) reported that the most important nematode associated with *Citrus* cultivation in Venezuela is *Tylenchulus semipenetrans*. This nematode was found associated with the cultivation of orange (*C. sinensis*) 'Valencia' grafted on tangerine 'Cleopatra' (*C. reshni*) rootstocks, lemon 'Volkameriano' (*C. volkameriana*) and citrange 'Carrizo' (*P. trifoliata* x *C. sinensis*) (Petit 1990a). Other nematodes present in soil samples, yet in lower densities, are those belonging to the genus *Helicotylenchus*, *Paratylenchus*, *Peltamigratus*, *Pratylenchus*, *Rotylenchus* (*R. reniformis*), *Trichodorus*, *Tylenchorhynchus* and *Tylenchus* (*T. costatus*) (Petit 1990b).

More recently, Crozzoli (2002) presented an excellent review with a very comprehensive list of phytoparasitic nematodes in Venezuela, responsible for the majority of the losses of the fruit industry in the country, including citrus.

Insects

Geraud (1983), after performing a comparative analysis between different pests of tomato and citrus, reports that during the 1950s–1960s, the most important insect pests of citrus plants in Venezuela were aphids (*Homoptera: Aphidae*) and the scales (*Homoptera: Diaspididae, Coccidae y Margarodidae*). Later, the mite causing the disease known as 'bronceado de los frutos', and identified as *Phyllocoptruta aleivora* (*Acarina: Eriophyidae*), was added to the list. However, it was only by the end of the 1960s that advances in entomological studies and direct technical assistance to growers allowed for an increase in knowledge of the main insects that affect citrus plants in the country. Not surprisingly, however, aphids and scales continued to be among the most important pests afflicting this crop.

During the years 1965–70, *Aleurocanthus woglumi* (Angeles *et al.* 1971; Geraud and Doreste 1977, cited by Geraud 1983; EPO 2003b), the citrus black fly, is reported for the first time in the most important citrus producing areas of Venezuela. It is worth recalling that by the end of the 1960s the use of insecticides was almost inexistent in citrus orchards, and that it was precisely the emergence of *A. woglumi*, as an important citrus pest, that stimulated its re-appearance. But on the other hand, it was also critical in

motivating studies on the natural enemies of *A. woglumi* (parasites, predators and pathogenic fungi) as potentially useful for establishing strategies of biological control (Ferrer 2001). It was a parasite wasp, *Prospaltella opulenta* (*Hymenoptera: Aphelinidae*) that allowed for successful and almost complete control of the citrus blackfly (Geraud *et al.* 1977, cited by Geraud 1983). In 1985, the Japanese white fly, *Parabemisia myricae* (Kuwana), was also reported as an important pest on different citrus cultivars (Chávez and Chávez 1985).

Interestingly, in 1976, Geraud reports for the first time in Venezuela the presence of the insect vector of CTV, the citrus black aphid, *T. citricida* in the region known as Mene Grande, Bolívar District, state of Zulia (Geraud 1976, cited by Geraud 1983).

Particularly in the Venezuelan Andes, in an excellent review, Briceño (2007) describes a series of insects that causes considerable damage on different species of *Citrus*. Among them, the causal agent of 'red scale', *Cryosomphalus aonidium*, is a cosmopolitan insect that infests any citrus plant under virtually any environmental condition and drastically reduces plant foliage resulting in decreased plant vigor and fruit quality. *Lepidosaphes beckii*, etiological agent of 'comma scale', attacks all parts of citrus trees, causes defoliation and produces yellowing around the scale. In the case of the 'white snow scale', or citrus 'queresa', caused by *Unaspis citri*, the damage is not as severe since it is also a target of a naturally occurring parasitism. Only the males feed on citrus plants during their immature stage. Therefore, it is only in old trees where layers of scales accumulate. The insect may impair tree respiration.

'Citrus moth', the leafminer *Phyllocnistis citrella*, was reported on oranges, tangerines and lemon in Venezuela for the first time in 1995, in Maturín, Monagas State and in the Andes in 1997. Damage is mainly caused by serpentine galleries, which in turn causes deformations and curling of the leaves. 'Orange moth', *Ecdyolopha* (*Gymnandrosoma*) *aurantiana*, on the other hand, is widely distributed in South America. In the Venezuelan Andes, it is mainly found on 'Naranja criolla' and late varieties. The lepidopteran perforates fruits, making them unmarketable and susceptible to attack by opportunistic fungi (Briceño 2005).

Among the aphids, *Aphis craccivora* has been reported to cause feeding damage on different species of citrus plants, while *A. gossypii* and *A. nerii* cause leaf deformation in *C. aurantium*. On the other hand, *A. spiraeicola*, although not an efficient virus vector, does cause severe deformations on diverse species of citrus plants. However, it is *Toxoptera citricida*, known as the 'citrus black aphid', that is the most damaging of the aphids affecting citrus crops in Venezuela. As an efficient vector of CTV, has led to the decimation of citrus plants in Venezuela killing over 6 million trees with the disease. This pest can cause severe deformations, when it is not viruliferous. Another species of the genus, *T. aurantii*, has also been reported in the country (Cermeli 2007).

To conclude with this section, it is important to emphasize the profound, positive impact that entomologists have caused in the citrus industry in Venezuela, and in fruitculture in general (Pacheco 2003, 2006). Aside from increasing our knowledge of important pests, many Venezuelan entomologists have ventured to use biological control from the very beginning of this scientific discipline in the country, with notable success. We have already mentioned the case of *A. woglumi*; however, prior examples have paved the way to biological control in Venezuela. During the years 1939–41, for example, C. H. Ballou (1941) introduced *Rodolia cardinalis*, a coccinellid, to control the cottony cushion scale, *Icerya purchasi* in citrus (Ferrer 2001). More recently, Linares *et al.* (2001) reported on the use of *Ageniaspis citricola* for the control of the aforementioned psyllid, *P. citrella*.

Others: weeds, algae and mollusks

Other pests can affect citrus production, without causing

economical impact comparable to that caused by viruses, fungi and nematodes. A good account of weeds present in citrus orchards, and other crops, is presented by Lárez (2007a, 2007b), but restricted to Monagas state, in eastern Venezuela. A comprehensive study of weeds in other areas of the country is not available. Regarding algae, there are episodic reports of the Chlorophyta, *Chepaleuros virescens* and *Trentepohlia* sp., but they are not considered major pests. Few reports also mention the parasitic plant *Langsdorfia* sp., which is not considered to be of relevance or economical importance.

Of the mollusks present in citrus orchards, the most prominent is *Orthalicus maracaibensis*, which can cause severe damage on the trunk and the cortex of young branches, leading to total loss of leaves (Fuentes 2006).

IMPROVEMENT AND PERSPECTIVES

It can be argued that research in Venezuela related to *Citrus* species in particular, and Rutaceae in general, has been scarce and devoid of continuity. Most of the studies regarding citrus have been oriented to horticultural management of the crop and limited to evaluations of the grafting potential rootstocks, and evaluation and control of diseases in commercial species and varieties (Laborem *et al.* 1993; MARNR 1996; Avilán *et al.* 1998; Laborem *et al.* 1998; Wagner *et al.* 1998; Cañizares and Rojas 2001; Ojasti 2001; Wagner *et al.* 2002; Quintero *et al.* 2004). Few ethnobotanical (Delascio 1985; Bermúdez and Velázquez 2002; Carrillo-Rosario and Moreno 2006), anatomical (Avilán *et al.* 1982, 1986; Laskowski 2006; Cañizares *et al.* 2007), germination (Leal 1970), phenological (Perez *et al.* 2004) and phytochemical studies have been carried out with citrus (Ojeda *et al.* 1998; Rincón *et al.* 2005). The more prominent findings have already been mentioned, or will be shown in the following paragraphs, but it is worth mentioning the persistence of a divorce between the research performed at universities and other governmental research institutes, and producer problems and needs. This fact, however, is not an exclusive feature attributable to citrus cultivation; neither is it a recent development. From 5,822 references of a Venezuelan agriculture bibliography (Badillo and Bonfanti 1957) only 14 were devoted to citrus (with eight of them only to citrus diseases). For the years ahead, matters did not change: the subjects of research on citrus are the same and research is still much needed, although the number of articles had increased steadily in the last 40 years. In the following lines a brief summary of several important findings on citrus plants in Venezuela in recent years are presented.

All edible species of the genus *Citrus* are very rich in ascorbic acid, flavonoids and essential volatile compounds, terpenoid oils in the parenchymatous tissues of leaves and branches (Rincón *et al.* 2005; Cañizares *et al.* 2007). They also contain coumarins of the bergapten group, which are frequently added to tanning preparations since they promote skin pigmentation. In Venezuela, Ojeda *et al.* (1998) identified a total of 51 chemical constituents of the essential oil from *C. limon*. The most abundant compound found was the monoterpene, limonene (65.65%), and the main aldehydes were geraniol (1.43%) and nerol (0.87%). Among the remaining hydrocarbons, they also identified β -pinene (11.0%) and γ -terpinene (9.01%) in elevated proportions. Besides, the authors reported on the presence of 3.79% of oxygenated compounds, of which 2.70% were composed of aldehydes, 0.53% alcohols and 0.56% esters. In another study, Rincón *et al.* (2005) focused on the composition and concentration of bioactive compounds of the citrus flour that is widely consumed in Venezuela (*C. sinensis*, *C. reticulata* and *C. paradisi*). High concentrations of ascorbic acid, magnesium, carotenoids, diet fibers and abundant polyphenolic compounds were reported. The latter showed high antioxidant efficiency, and since they are more abundant in tangerine peels, they are recommended for the prevention of cardiovascular diseases and other health threat-

ening conditions related to lipid oxidation.

One of the first and persistent systematic attempts made to improve citrus cultivation and productivity was a study evaluating different rootstocks for pathogen resistance (Rondón *et al.* 1993) and tolerance to diverse sources of abiotic stress. Regarding planting materials, micrografting was proven to be valuable in the generation of germplasm free from exocortis and psorosis (Monteverde *et al.* 1986), and yet no attempts have been made to produce citrus plantlets free from virus and viroids on a greater scale. In the same vein, in a study on pollen germination, Leal (1970) concluded that *P. trifoliata* and some oranges and tangelos (*C. reticulata* x *C. paradisi*) are appropriate germplasm for improvement programs. However, there is no evidence that these findings have ever been put in practice.

In a technical report on the use of biotechnology in Venezuela, García (2005) mentions the application of clonal micropropagation and the production of clean seeds, the advances in the use of molecular markers to citrus and their pathogens. She also touted an increase in the number of scientists in the country with proficient skills in molecular biology and predicts an increase in the technological input to citriculture in Venezuela. For example, the detection of viruses by multiplex polymerase chain reaction (Roy *et al.* 2005) has been proven an effectiveness method of detecting CTV, which remains the prevalent viral disease in citrus in Venezuelan orchards (unpublished). On the other hand, the control of *Papaya ringspot virus* by a transgenic approach (Fermin *et al.* 2004), set an example for the application of modern techniques for the control important pathogens in fruit crops, but the country does not appear to be ready for the acceptance and commercialization of genetically modified crop plants (Fermin *et al.* 2005).

Few attempts to improve *Citrus* germplasm by Venezuelan researchers using *in vitro* technologies in hybrids have been applied, in some cases with notable success (Viloria *et al.* 2005). This valuable effort represents another example of scientists already prepared to assume the challenge of modernizing Venezuelan citriculture since, although *in vitro* tissue culture techniques have been widely used in the country, citrus species might be considered so far a neglected target in this regard.

In 2007, the Interamerican Web on Citrus held a meeting in Venezuela, where members of three different countries, as well as producers, gathered to discuss pressing issues related to citrus cultivation. It is expected that the reactivation of the working group might lead to a concerted effort to assist producers in designing a modernized strategy that might help increase citrus production. Some governmental efforts aimed at providing technical and financial assistance to citrus growers are under way in different parts of the country, but it is still too early to predict the final outcome of what seems to be a serious attempt to recover citrus production.

CONCLUDING REMARKS

The citrus industry in Venezuela, including production and processing, seems to be recovering from a decade of low productivity, low technical input, scarce research and poor financial support. Additionally, there are certain aspects of the Venezuelan society, as a whole, that provide an optimistic view for the future of the citrus industry: First, a well formed group of phytopathologists showing a renewed interest in the main pathogens that attack fruit trees, including the diverse species and varieties of *Citrus* cultivated in the country; second, by all measures, the economic situation of the vast majority of the inhabitants of the country has improved, which has led, and will continue doing so, to an increase in food availability and consumption, including fruits; third, although still very low, exports of citrus might stimulate an increase in production for both local and external markets. However, the country needs a renewal of its citrus genetic material. The main obstacle for a rational, comprehensive and realistic genetic improvement program

is the absence of previous experience in this field for this particular important crop. To our knowledge, the limited citrus germplasm in the country (Pérez *et al.* 1998; Rincón *et al.* 2006) is not sufficient to support an improvement program (for example, there are only 93 entries for citrus species at CENIAP, according to Pérez *et al.* (1998), of which less than 15 have been characterized), nor is the almost inexistent experience in tissue culture or transformation of citrus species enough to sustain an improvement program adapted to modern biotechnology.

REFERENCES

- Angeles N, Dedordy J, Paredes P, Requena J (1971) Mosca prieta (*Aleurocanthus woglumi* Ashby) de los cítricos en Venezuela. *Agronomía Tropical* 21, 71-75
- Arnal E, Russell LM, Debrot E, Ramos F, Cermeli M, Marciano R, Montagne A (1993) Lista de moscas blancas (Homoptera:Aleyrodidae) y sus plantas hospederas en Venezuela. *The Florida Entomologist* 76, 365-381
- Avilán L, Dorantes I, Ruiz J, Rodríguez M (1998) Descripción de las limas y limones de la colección del Centro Nacional de Investigaciones Agropecuarias. *Agronomía Tropical* 48, 41-52
- Avilán L, Leal F, Meneses L (1982) Distribución del sistema radical en las combinaciones de naranjo dulce (*Citrus sinensis*) y grapefruit (*Citrus paradisi*) sobre patrones cajero (*Citrus aurantium*) en suelos calcáreos de la hoya del Lago de Valencia (Venezuela). *Agronomía Tropical* 32, 155-170
- Avilán L, Ruiz J (1999) 70 toneladas de naranjas por hectárea en la citricultura nacional. ¡Una realidad! *FONALAP Divulga* 61, 4 pp
- Avilán L, Velarde C, Meneses L (1986) Distribución del sistema radical de los patrones de cítricos naranjo agrio (*Citrus aurantium* L.), cleopatra (*C. reshni* Hort. ex Tan.) y volkameriana (*C. volkameriana* Pasq.). *Agronomía Tropical* 36, 97-113
- Badillo VM, Bonfanti C (1957) Índice bibliográfico agrícola de Venezuela. Fundación Eugenio Mendoza. Caracas, 305 pp
- Ballou C (1941) Notas sobre la introducción de un predador para el control de la escama algodonosa del naranjo. *Boletín de la Sociedad Venezolana de Ciencias Naturales* VII, 121-122
- Benaccio S, Cañizales R, Avilán W (1985) Zonificación agroecológica del cultivo de la naranja (*Citrus sinensis* L.) en Venezuela. FONALAP, Barquisimeto, Venezuela. Editora Boscan, 49 pp
- Bernúdez A, Velázquez D (2002) Etnobotánica médica de una comunidad campesina del estado Trujillo, Venezuela: un estudio preliminar usando técnicas cuantitativas. *Revista de la Facultad de Farmacia* 44, 2-6
- Briceño A (2005) Plagas chupadoras de las cítricas (naranjo, limonero, mandarina) en Los Andes venezolanos. *Boletín Divulgativo IIAP (Universidad de Los Andes, Mérida, Venezuela)* 60, 20-24
- Briceño A (2007) Importancia agrícola de los principales insectos plagas en Los Andes Venezolanos. Consejo de Estudios de Postgrado, Facultad de Ciencias Forestales y Ambientales, Universidad de Los Andes. Mérida, Venezuela, 387 pp
- Caballero R (1995) La etnobotánica en las comunidades negras e indígenas del Delta del río Patía. Colección Biblioteca 'Abya-Yala' (Vol 26). Universidad Nacional de Colombia, Medellín, 248 pp
- Cañizales A, Rojas E (2001) Efecto de la cianamida de hidrógeno y el ácido 2-cloroetil fosfónico sobre la foliación, floración y fructificación de la lima 'Tahiti'. *Bioagro* 13, 10-14
- Cañizales AE, Salcedo F (1994) Evaluación del desarrollo vegetativo de variedades de limones (*Citrus limon*) en suelos de sabanas del Estado Monagas. V Congreso Nacional de Frutales. Maracay, Venezuela, 124 pp
- Cañizales A, Sanabria M, Rojas E (2007) Anatomía del tallo de Lima Tahiti (*Citrus latifolia* Tanaka). *Revista UDO Agrícola* 7, 221-227
- Carrillo-Rosario T, Moreno G (2006) Importancia de las plantas medicinales en el autocuidado de la salud en tres caseríos de Santa Ana Trujillo, Venezuela. *Revista de la Facultad de Farmacia* 48, 21-28
- Cedeño L, Palacios-Prú E (1992) Identificación de *Botryodiplodia theobromae* como la causa de lesiones y gomosis en cítricos. *Fitopatología Venezolana* 5, 10-13
- Cermeli M (2007) Áfidos de importancia agrícola en Venezuela. In: *Plagas Agrícolas de Venezuela*. Monografía, Sociedad Venezolana de Entomología, Maracay, 29 pp
- Cermeli M, Morales P, Godoy F (2000) Presencia del psílido asiático de los cítricos *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) en Venezuela. *Boletín de Entomología Venezolana* 15, 235-243
- Chardon CE, Toro RA (1934) Mycological explorations of Venezuela. Monographs of the University of Puerto Rico, Series B, Number 2. The University of Puerto Rico, Rio Piedras, 353 pp, 33 plates and a map
- Chávez H, Chávez A (1985) *Parabemisia myricae* (Kuwana) Homoptera: Aleyrodidae, plaga potencial en cítricos y aguacate en Venezuela. *Boletín de Entomología Venezolana N.S.* 4, 71-72
- Ciferri R (1950) A decline of citrus plants in Venezuela. *Nature* 165, 32
- Clawson DL, Raymond EC (1982) Evolution of land-use patterns and agricultural systems. *Mountain Research and Development* 2, 265-272
- CONAPLAMED: Comisión Nacional para el Aprovechamiento de Plantas Medicinales (2000) *Cuaderno de fitoterapia clínica*. Edigráfica, Mérida, 172 pp
- Contreras N, Rondón A (1980) Etiología de la mancha parda de los cítricos en Venezuela. *Agronomía Tropical* 35, 111-115
- Crozzoli R (2002) Especies de nemátodos fitoparasíticos en Venezuela. *Inter-ciencia* 27, 345-364
- Cunill P, Pino E, Rodríguez M, Rodríguez JA, Romero P, Sanoja M, Vargas I (2007) La Geografía Histórica del poblamiento territorial venezolano (In *GeoVenezuela*, Vol I). Fundación Empresas Polar, Caracas, 416 pp
- Delascio F (1985) *Algunas plantas usadas en la medicina empírica venezolana*. Dirección de Investigaciones Biológicas, División de Vegetación Jardín Botánico. INPARQUES, Caracas, 186 pp
- Dennis RWG (1970) Kew Bulletin Additional Series III. Fungus flora of Venezuela and adjacent countries. Verlag von J. Cramer, 531 pp
- Díaz C, Salas G (1973) Lista de patógenos de las plantas cultivadas en Venezuela. Sociedad Venezolana de Fitopatología. Boletín Especial N° 2 (Monograph), 59 pp
- EPP0 (2003a) Data Sheets on Quarantine Pests: *Spiroplasma citri*. Prepared by CABI and EPP0 for the EU under Contract 90/399003, 7 pp
- EPP0 (2003b) Data Sheets on Quarantine Pests: *Aleurocanthus woglumi*. Prepared by CABI and EPP0 for the EU under Contract 90/399003, 5 pp
- EPP0 (2004) Data Sheets on Quarantine Pests: Citrus blight disease. Prepared by CABI and EPP0 for the EU under Contract 90/399003, 6 pp
- EPP0 (2007) Data Sheets on Quarantine Pests: Citrus leprosis 'rhabdovirus'. Prepared by CABI and EPP0 for the EU under Contract 90/399003, 6 pp
- Erwin DC, Ribeiro OK (1996) *Phytophthora Diseases Worldwide*, APS Press, St. Paul, Minnesota, 562 pp
- Farr DF, Rossman AY, Palm ME, McCray EB (2008) Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. Available online: <http://nt.ars-grin.gov/fungaldatabases/>
- Ferguson JJ (2002) Medical uses of *Citrus*. Horticultural Sciences Department, Florida. Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, 3 pp
- Fermin G, Ingleis V, Garboza C, Rangel S, Dagert, M, Gonsalves D (2004) Engineered against *Papaya ringspot virus* in Venezuelan transgenic papayas. *Plant Disease* 88, 516-522
- Fermin G, Tennant P, Gonsalves C, Lee D, Gonsalves D (2005) Comparative development and impact of transgenic papayas in Hawaii, Jamaica, and Venezuela. In: Peña L (Ed) *Transgenic Plants: Methods and Protocols. Methods in Molecular Biology* (Vol 286), Walker JM (Series Ed), Humana Press, Totowa, New Jersey, pp 399-430
- Fernández F, Rosales CJ, Kern F, Szumkowski W, Whitcomb WH, Labrador JR, Fernández A, Doreste E (1957) Lista preliminar de nombres comunes de algunos insectos dañinos de Venezuela. Monografía, Facultad de Agronomía, Universidad Central de Venezuela, Maracay, Venezuela, 14 pp
- Ferrer F (2001) Biological control of agricultural insect pests in Venezuela: advances, achievements, and future perspectives. *Biocontrol/News and Information* 22, 67N-74N
- Fuentes L (2006) Moluscos de importancia agrícola. *CENIAP Hoy* 11, 10 pp
- Fundación Shell (1966) *Servicio Shell para el Agricultor: Cítricas* (Serie A, N° 27) (2nd Edn), Editorial Nuestra América, Cagua, Venezuela, 82 pp
- García E (2005) Diagnóstico sobre la situación actual de la biotecnología y bioseguridad en la República Bolivariana de Venezuela. Monografía, Oficina Nacional de Diversidad Biológica, Ministerio del Poder Popular para el Ambiente, Caracas, 111 pp
- Garnsey SM, Chagas CM, Chiavegato LG (1988) Leprosis and zonate chlorosis. In: Whiteside JO, Garnsey SM, Timmer LW (Eds) *Compendium of Citrus Diseases*, APS Press, The American Phytopathological Society, St. Paul, Minn., pp 43-44
- Geraud F (1983) Plagas de Tomate y en cítricos y su control por insecticidas en Venezuela. Un análisis comparativo. *Revista de la Facultad de Agronomía* 6, 758-767
- Gianessi LP, Silvers CS, Sankula S, Carpenter JE (2002) Viral resistant citrus. In: *Plant Biotechnology: Current and Potential Impact for Improving Pest Management in U.S. Agriculture. An Analysis of 40 Case Studies*, National Center for Food and Agricultural Policy, 10 pp
- Gil R, Carmona J (2003) *Herbolario Tradicional Venezolano*, Universidad de Los Andes. Consejo de Publicaciones, Mérida, 218 pp
- Gmitter FG, Hu X (1990) The possible role of Yunnan, China, in the origin of contemporary *Citrus* species (Rutaceae). *Economic Botany* 44, 267-277
- Halbert SE, Nuñez CA (2004) Distribution of the Asian citrus psyllid, *Diaphorina citri* Kuwayama (Rhynchota: Psyllidae) in the Caribbean basin. *Florida Entomologist* 87, 401-402
- Hermoso L, Escala M (2002) Anatomía foliar de *Pilocarpus goudotianus* Tul. (Rutaceae). *Caldasia* 24, 269-275
- Hernández L, Ochoa F, Derrick K, Barreta MJ, Castejón O, Chacín F, Machado TW (1993) Relación entre *Xylella fastidiosa* y el decaimiento repentino de los cítricos. *Fitopatología Venezolana* 6, 42-43
- Hilf ME, Mavrodieva VA, Garnsey SM (2005) Genetic marker analysis of a global collection of isolates of *Citrus tristeza virus*: characterization and distribution of CTV genotypes and association with symptoms. *Phytopathology*

- 95, 909-917
- Hokche O, Berry PE, Huber O** (2008) Nuevo catálogo de la flora vascular de Venezuela. Fundación Instituto Botánico de Venezuela Dr. Tobias Lasser, Caracas, 859 pp
- Hoyos J** (1994) *Frutales en Venezuela* (2nd Edn), Sociedad de Ciencias Naturales La Salle, Caracas, Venezuela, 381 pp
- Hurtado MA, Hill Kim R** (1990) Seasonality in a foraging society: variation in diet, work effort, fertility, and sexual division of labor among the Hiwi of Venezuela. *Journal of Anthropological Research* **46**, 293-346
- Játem A, Ricardí M, Adamo G** (1997) Herbal traditional medicine of Venezuelan Andes: an ethnopharmacological study. *Phytotherapy Research, Special Issue: Proceedings of the Second Internacional Symposium on Natural Drugs* **12 S2**, S53-S59
- Jones E, Ocanto L, Guevara J** (1964) Bailadores: An agro-social study of a rural Venezuelan region. Instituto Caribe de Antropología y Sociología, Caracas, 192 pp
- Judd WS, Campbell CS, Kellogg EA, Stevens PF, Donoghue MJ** (2002) *Plant Systematics: A Phylogenetic Approach* (2nd Edn), Sinauer Associates, Inc. Sunderland, 576 pp
- Junta del Acuerdo de Cartagena** (1996) Resolución 419: Inventario subregional de plagas y enfermedades de los vegetales de importancia económica para el área Andina. *Gaceta Oficial del Acuerdo de Cartagena* **223**, 1-115
- Keshava B** (1985) *Herbolario tropical*, Editorial Texto, Caracas, Venezuela, 176 pp
- Knorr LC, Denmark HA** (1970) Injury to citrus by the mite *Brevipalpus phoenicis*. *Journal of Economic Entomology* **63**, 1996-1998
- Kyprianou M** (2006) Commission decision of 5 July 2006 recognizing certain third countries and certain areas of third countries as being free from *Xanthomonas campestris* (all strains pathogenic to Citrus), *Cercospora angulensis* Carv. et Mendes and *Guignardia citricarpa* Kieley (all strains pathogenic to Citrus), (notified under document number C(2006) 3024) (2006/000/EC). *Official Journal of the European Union*, L 187/35:35-36
- Laborem G, Monteverde E, Rangel L, Espinoza M** (1998) Evaluación de la calidad de los frutos de naranjos, mandarinos, pomelos e híbridos injertados sobre 'Cleopatra'. *Bioagro* **10**, 35-39
- Laborem G, Rangel L, Espinoza M** (1996) Cítricos exportables y su manejo. *FONALAP Divulga* **52**, 8-11
- Laborem G, Reyes F, Rangel L** (1993) Calidad de la cosecha de la naranja 'Valencia' sobre ocho patrones. FONALAP - CENIAP, Instituto de Investigaciones Agronómicas, Serie A N° 10, Maracay, Venezuela, 32 pp
- Lans C** (2007) Ethnomedicines used in Trinidad and Tobago for reproductive problems. *Journal of Ethnobiology and Ethnomedicine* **3**, 13-24
- Lans CA** (2006) Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus. *Journal of Ethnobiology and Ethnomedicine* **2**, 45-55
- Lárez A** (2004) Las plantas medicinales en el municipio Caripe, Estado Monagas. Universidad de Oriente, Monagas, Venezuela, 106 pp
- Lárez A** (2007a) Claves para identificar malezas asociadas con diversos cultivos en el estado Monagas, Venezuela. I. Monocotiledóneas. *Revista UDO Agrícola* **7**, 79-90
- Lárez A** (2007b) Claves para identificar malezas asociadas con diversos cultivos en el estado Monagas, Venezuela. II. Dicotiledóneas. *Revista UDO Agrícola* **7**, 91-121
- Laskowski L** (2006) Características de la abscisión del fruto de naranja *Citrus sinensis* (L.) Osbeck var. salustiana. *Bioagro* **18**, 1316-13361
- Leal F, Trujillo A** (1970) Patrones para cítricos. *Noticias Agrícolas (Venezuela)* **9**, 131-132
- Leal F** (1970) Germinación del polen en *Citrus* y *Poncirus*. *Agronomía Tropical* **20**, 323-329
- Linares B, Hernández J, Morillo J, Hernández L** (2001) Introducción de *Agynaspis citricola* Logvinovskaya, 1983 (Hymenoptera: Encyrtidae) para el control del 'minador de la hoja de los cítricos' *Phyllocnistis citrella* Staiton (Lepidoptera: Gracillariidae: Phyllocnistinae) en el estado Yaracuy, Venezuela. *Entomotropica* **16**, 143-145
- Mabberley DJ** (2000) A classification for edible *Citrus* (Rutaceae). *Telopea* **7**, 167-172
- Machado-Allison C** (2007) *Consumo de Alimentos en Venezuela*, Ediciones IESA, Caracas, 272 pp
- MARNR: Ministerio del Ambiente y de los Recursos Naturales Renovables** (1996) Venezuela: informe nacional para la conferencia técnica internacional de la FAO sobre los recursos fitogenéticos. Monografía. Caracas, Venezuela, 117 pp
- Martínez J, Sulbarán de Ferrer B, Ojeda de Rodríguez G, Ferrer A, Nava R** (2003) Actividad antibacteriana del aceite esencial de mandarina. *Revista de la Facultad de Agronomía-LUZ* **20**, 502-512
- Milliken W, Albert B** (1997) The use of medicinal plants by the Yanomami Indians of Brazil, part II. *Economic Botany* **51**, 264-278
- Mendí R** (1992) History of CTV in Venezuela. In: Lastra R, Ochoa FL, Lee RF, Niblett CL, Rocha-Peña MR, Gamsey SM, Yokomi RK (Eds) *Citrus tristeza virus and Toxoptera citricidius* in Central America: Development of Management Strategies and the Use of Biotechnology for Control. CATIE-University of Florida-INIFAP/SAH-Universidad Central de Venezuela-USDA. Maracay, Venezuela, pp 137-141
- Ministerio del Poder Popular para la Salud** (2007) Hojas de balance de alimentos, 2002-2004. Monografía, Instituto Nacional de Nutrición, Caracas, Venezuela, 333 pp
- Monteverde E** (1983) La tristeza de los cítricos en Venezuela. *FONALAP Divulga* **11**, 7 pp
- Monteverde E, Espinoza M, Ruiz J** (1981) Presence of the virus exocortis in three citrus orchards of Carabobo high valleys (Venezuela). Convención Nacional de Fruticultores, Maracaibo, Venezuela, 7 pp
- Monteverde E, Espinoza M, Ruiz J, Briceño M** (1992) Evaluación de psorosis-concave gum, exocortis y cachexia-xyloporosis en árboles de naranjo dulce en los valles altos de Carabobo, Venezuela. *Agronomía Tropical* **42**, 137-150
- Monteverde E, Espinoza M, Ruiz J** (1985) Identificada la xyloporosis de los cítricos en Venezuela con plantas indicadoras. *Agronomía Tropical* **35**, 173-176
- Monteverde EE, García ML, Briceño M** (1986) Obtención de plantas cítricas libres de psorosis y exocortis en árboles infectados a través de la microinjertación de ápices *in vitro*. *Agronomía Tropical* **36**, 5-14
- Monteverde E, Laborem G, Ruiz J, Espinoza M, Guerra C** (1996) Evaluación del naranjo Valencia sobre siete patrones en los valles altos Carabobo-Yaracuy, Venezuela 1984-1991. *Agronomía Tropical* **46**, 371-393
- Moore GA** (2001) Oranges and lemons: clues to the taxonomy of *Citrus* from molecular markers. *Trends in Genetics* **17**, 536-540
- Mora EM, Rojas J** (2007) Los cultivos líderes de la agricultura venezolana. *Agroalimentaria* **25**, 33-44
- Moreno MJ, Belén DR, García D, Mendoza L** (2006) Evaluación del contenido de carotenoides totales en cáscaras de algunas variedades de naranjas venezolanas. *Revista de la Facultad de Agronomía-LUZ* **23**, 301-309
- Müller AS** (1940) Observaciones sobre las enfermedades de las plantas cultivadas en Venezuela, 1937-1940. Monografía, Estación Experimental de Agricultura y Zootecnia, Caracas, Venezuela, 18 pp
- Ochoa F, Carballo O, Trujillo G, Mayoral M, Lee R** (1993) Biological characterization and evaluation of cross protection potential of *Citrus tristeza virus* isolates in Venezuela. In: Moreno P, Da Graça JV, Timmer LW (Eds) *Proceedings of the 12th Conference of the International Organization of Citrus Virologists*, Riverside, CA, p 1-7
- Ojasti J** (2001) *Estudio Sobre el Estado Actual de las Especies Exóticas: Estudio Nacional*, Biblioteca Digital Andina, Secretaría General de la Comunidad Andina, Caracas, 223 pp
- Ojeda G, Morales V, González N, Cabrera L, Sulbarán B** (1998) Composition of Venezuelan lemon essential oil *Citrus limon* (L.) Burm.f. *Revista de la Facultad de Agronomía* **15**, 343-349
- Ordosgoitty A, Marquez O, Garces N** (1971) Bibliografía fitopatológica Venezolana. Centro de Investigaciones Agronómicas, Maracay, Venezuela. *Boletín Bibliográfico* **7**, 68 pp
- Ortega E** (1986) Métodos de detección virus y viroides en cítricos. *FONALAP Divulga* **22**, 4 pp
- Pacheco G** (2003) La institucionalización de la investigación agrícola en Venezuela: los primeros tiempos, 1870-1935. *Agroalimentaria* **16**, 65-82
- Pacheco G** (2006) La construcción de la entomología agrícola en Venezuela y la Facultad de Agronomía de la UCV: una visión histórica. *Bitácora-e* **2**, 45-68
- Patiño VM** (1963) Plantas cultivadas y animales domésticos en América equinoccial. Tomo I Frutales. Imprenta Departamental, Cali, Colombia, 547 pp
- Patiño VM** (1969) *Plantas Cultivadas y Animales Domésticos en América Equinoccial. Tomo IV: Plantas Introducidas*, Imprenta Departamental, Cali, Colombia, 573 pp
- Patiño VM** (1982) Biotic resources for potential development. *Mountain Research and Development* **2**, 333-336
- Peréz D, Gutiérrez M, Mazzani E, Barreto T, Segovia V, Marín CR** (1998) Recursos fitogenéticos en Venezuela. Fondo Nacional de Investigaciones Agropecuarias. Centro Nacional de Investigaciones Agropecuarias (Serie C. N° 42), Maracay, Venezuela, 93 pp
- Pérez J, Lago M, Mata O** (1975) Nota sobre el uso de la harina de cítricos en dietas para pollos barrileros. *Agronomía Tropical* **26**, 261-268
- Pérez M, Soto E, Avilán L** (2004) Descripción de la fenología en tres cultivares de cítricos en la zona central de Venezuela. *Revista de la Facultad de Agronomía* **21**, 102-108
- Petit P** (1990a) Reconocimiento de nemátodos fitoparásitos asociados a frutas de importancia económica en Venezuela. *Fitopatología Venezolana* **3**, 2-5
- Petit P** (1990b) Presencia del nemátodo de las cítricas (*Tylenchulus semipenetrans*) en la zona citrícola del centro del Venezuela. *Fitopatología Venezolana* **4**, 11-12
- Pretorius MC, Crous PW, Groenewald JZ, Braun U** (2003) Phylogeny of some cercosporoid fungi from *Citrus*. *Sydowia* **55**, 286-305
- Quijada O, Jiménez O, Mathes M, Monteverde E** (2002) Evaluación del limero Tahiti sobre 10 portainjertos en la planicies de Maracaibo. *Revista de la Facultad de Agronomía* **19**, 173-184
- Quiñero I, Alvarez R, Zambrano J, Materano W, Maffei M** (2004) Evaluación de plantaciones de naranjo 'Valencia' a diferentes pisos altitudinales, en Trujillo-Venezuela. *Revista de la Facultad de Agronomía* **21**, 277-284
- Rangel E, Cuello R, Ascanio M, Centeno F, Ruiz J** (2000) Incidencia del virus de la tristeza de los cítricos en algunas localidades de los estados Ara-

- gua, Carabobo y Yaracuy, Venezuela. *Fitopatología Venezolana* **13**, 19-21
- Reyes F, Ruiz J** (1984) Comportamiento en vivero de patrones cítricos tolerantes a tristeza. *Agronomía Tropical* **34**, 35-41
- Reynel C** (1995) New Andean *Zanthoxylum* (Rutaceae) with distinctive vegetative characters. *Novon* **5**, 362-367
- Ricardi MH** (1991) *Familias de Dicotiledóneas Venezolanas* (Vol II), Universidad de Los Andes, Mérida, Venezuela, 192 pp
- Rincón A, Pérez D, Romero S** (2006) Agricultura tropical sustentable y biodiversidad. *CENIAP Hoy* **11**, 16 pp
- Rincón AM, Vásquez AM, Padilla FC** (2005) Composición química y compuestos bioactivos de las harinas de cáscaras de naranja (*Citrus sinensis*), mandarina (*Citrus reticulata*) y toronja (*Citrus paradisi*) cultivadas en Venezuela. *Archivos Latino Americanos de Nutrición* **55**, 1-6
- Rivas B, Yanez V, Carrizales L, Sanchez M** (2006) Hongos asociados a la caída prematura de frutos en lima persa (*Citrus latifolia* Tan) y evaluación de su control químico. *Bioagro* **18**, 31-39
- Rojas E, Scorza JV** (1991) The use of lemon essential oil as a sandfly repellent. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **85**, 803
- Rondón A** (1990) Enfermedades de los frutales en Venezuela. Instituto de Investigaciones Agronómicas. FONAIAP - CENIAP, Serie B N° 9, Maracay, Venezuela, 96 pp
- Rondón A** (1992) Enfermedades Fungosas de los Cítricos. *FONAIAP Divulga* **39**, 4 pp
- Rondón A, Angeles N, Leal F** (1970) La 'petrificación' de los cítricos en Venezuela. *Agronomía Tropical* **20**, 131-134
- Rondón A, Hung G, Reyes F, Solórzano R** (1993) Reacción de patrones cítricos a *Phytophthora nicotianae* B. de Haan var. *parasitica* (Dastur) Waterh, en condiciones de umbráculo. *Agronomía Tropical* **43**, 117-125
- Roy A, Fayad A, Barthe G, Brlansky RH** (2005) A multiplex polymerase chain reaction method for reliable, sensitive and simultaneous detection of multiple viruses in citrus trees. *Journal of Virological Methods* **129**, 47-55
- Rudolph WE** (1944) Agricultural possibilities in northwestern Venezuela. *Geographical Review* **34**, 36-56
- Scora RW** (1975) On the history and origin of Citrus. *Bulletin of the Torrey Botanical Club* **102**, 369-375
- Serpa D** (1965) Nomenclatura y clasificación de las cítricas en Venezuela. *Revista de la Facultad de Agronomía* **3**, 5-8
- Spaulding P** (1961) Foreign diseases of forest trees of the world. U.S.D.A. *Agriculture Handbook* **197**, 1-361
- Tamasaulas R, Ruiz H, Theis W, de Basilio V, Herrera J, Aguirre A, Roa N** (1997) Nuevo método de control de la coccidiosis aviar: Pruebas *in vitro* y de eficacia en piso con pollos de engorde para la evaluación de un desinfectante a base de extractos de cítricos en Venezuela. *Archivo Latinoamericano de Producción Animal* **5**, 612-615
- Taylor MS** (1989) Plant taxa described by Julian A. Steyermark. *Annals of the Missouri Botanical Garden* **76**, 652-780
- Urtiaga R** (1986) *Índice de Enfermedades en Plantas de Venezuela y Cuba*, Impresos Nuevo Siglo. S.R.L., Barquisimeto, Venezuela, 202 pp
- Valbuena M** (1996) Evaluación del limón Volkameriano (*Citrus volkameriano* Pasq.) y mandarina Cleopatra (*Citrus reshii* Hort.) como patrones de la lima Persa (*Citrus latifolia* Tam.) en la cuenca media del río Guasare, Sierra de Perijá, Estado Zulia, Venezuela. *Revista de la Facultad de Agronomía* **13**, 139-151
- Vila MA** (1981) *Plantas de Cultivo y de Recolección en la Geohistoria Venezolana*, Ediciones de la Facultad de Humanidades y Educación, Escuela de Geografía, Universidad Central de Venezuela, Caracas, 429 pp
- Viloria Z, Grosser JW, Bracho B** (2005) Immature embryo rescue, culture and seedling development of acid citrus fruit derived from interloid hybridization. *Plant Cell, Tissue and Organ Culture* **82**, 159-167
- Vit P, Hernández P, Mercado R** (2006) Revisión sobre el conocimiento de las mieles uniflorales venezolanas. *Revista de la Facultad de Medicina, Universidad de Los Andes* **15**, 29-39
- Wagner M, Laborem G, María C, Medina G, Rangel L** (2002) Efecto de diferentes patrones de cítricas e intervalos de riego sobre la calidad y producción de la naranja 'Valencia'. *Bioagro* **14**, 71-76
- Wagner M, Laborem G, Medina G, Rangel L** (1998) Efecto del patrón y la frecuencia de riego sobre el nivel foliar de prolina en el naranjo 'Valencia'. *Bioagro* **10**, 76-79
- Yepez G, Meredith JA** (1970) Nematodos fitoparásitos en cultivos de Venezuela. *Revista de la Facultad de Agronomía* **5**, 3-8